

Environmental Factors

Nutrients


K. Raja Reddy

Krreddy@pss.msstate.edu



Mississippi State University
A Land-Grant Institution

Environmental and Cultural Factors Limiting Potential Yields

- 
- Atmospheric Carbon Dioxide
 - Temperature (Extremes)
 - Solar Radiation
 - Water
 - Wind
 - **Nutrients (N and K)**
 - Others, ozone etc.,
 - Growth Regulators (PIX)

Nutrients - Objectives


The objectives of this lecture are to:

- Learn temporal trends in fertilizer usage (Major nutrients).
- Influence of major nutrients on plant growth and development.

Major Nutrients


Trends and some Statistics

Trends in World Commercial Fertilizer Use




Trends in U.S. Commercial Fertilizer Use

(China, USA and India)



Trends in U.S. Commercial Fertilizer Use

(Nitrogen, Potash and Phosphate)




Trends in U.S. Total Commercial Fertilizer Use

(Primary, Secondary and Micronutrients)

Commercial fertilizer use depends on variety of factors:

- Soil
- Climate and weather
- Feasible technology
- Crop mix
- Crop rotations
- Technological change
- Govt. programs
- Commodity and fertilizer prices
- Affordability



Major Nutrients and Their Influences

Nutrient Supply and Plant Growth




Fig. 12.1 Relationship between nutrient supply and growth.

Nitrogen Supply and Plant Growth




Fig. 8.16 Schematic representation of the effect of increasing levels of nitrogen supply on the root and shoot growth of cereal plants.

Nitrogen and Crop Yield





Figure 5.25 Response curve showing diminishing increments.




Question:

- Do processes within a crop vary in their response to nutrients?


Nitrogen and Cotton Growth and Development

Leaf developmental response to N and elevated CO₂





Nitrogen and Crop Growth and Development

Leaf growth response to N and elevated CO₂




Cotton leaf area at leaf unfolding



Caster mature leaf sizes


✓N treatments were imposed when leaf 5 was just unfolding



Nitrogen and Crop Growth and Development


Cotton leaf growth response to N and elevated CO₂

RLER = Relative Leaf Expansion Rate




Nitrogen and Cotton Growth and Development

Stem elongation response to N and elevated CO₂




Nitrogen and Cotton Growth and Development

Stem Elongation Rate Response to N and elevated CO₂




Nitrogen and Cotton Growth and Development

Leaf photosynthetic response to N and elevated CO₂




Nitrogen and Cotton Growth and Development

Relative Rates of Photosynthesis



Can we use one function for all processes in a given crop?

Functional relationships – cotton for growth and developmental processes




Questions:

- Do species vary in their response to nutrients?
- How about functional groups such as C_3 versus C_4 ?
- Is there a difference between the functional groups in their response to nutrients?


Can we apply cotton algorithms for other crops?

N and Photosynthesis – Functional Groups




Can we apply cotton algorithms for other crops?

N and Photosynthesis – Several Crops




Can we apply cotton algorithms for other crops?

N and Photosynthesis – Several Crops




Photosynthesis - Variability Among Species

Response to Leaf Nitrogen




Can we apply cotton algorithms for other crops?

N and several crops – Stem elongation rates



Can we apply cotton algorithms for other crops?

N and several crops – Leaf area expansion rates




Summary and Conclusions

Nitrogen Responses across Species and Processes


- Functional algorithms varied among crop species and even among crop species within a functional physiological group such as C₃ or C₄ species.
- Functional algorithms varied among crop processes for a given species.
- Among the growth, developmental and physiological processes, leaf growth was more responsive to leaf N than other processes in almost all crops.
- N also affects cell division and cell elongation process leading to a cascade of effects on several processes in plants, and finally yield.

Potassium Supply and Plant Growth



7 Effect of potassium supply on grain yield of wetland rice and incidence of stem rot (*thosporium sigmoideum*). Basal dressing of nitrogen and phosphorus constant at 120 and 60 kg ha⁻¹, respectively. (Based on Ismunadji, 1976.)

Potassium Supply and Plant Growth



8 Severity of leaf spot disease (*Helminthosporium cynodontis*) and dry matter yield in bermudagrass (*Cynodon dactylon* L. Pers.) versus leaf potassium content. (Reproduced in Matocha and Smith, 1980, by permission of the American Society of Agronomy.)

Potassium – Cotton Growth

<3.05



1.15



0.94



0.39




0.30




Visual Symptoms
and leaf K, %

Potassium and Cotton Growth and Development



Photosynthesis and Environment

Response to phosphorus – Sub-to supra-optimal supply of Pi



Summary and Conclusions

Nutrient Responses across Species and Processes

- Functional algorithms or responses varied among crop species.
- Functional algorithms varied among crop processes for a given species.
- Similar to N effects, among the growth, developmental and physiological processes, leaf growth was more responsive to leaf K.
- The effects of P on various processes are less quantified to arrive a conclusion.