

Crop Growth and Phenology — Species Variability

K. Raja Reddy
Mississippi State University
Mississippi State, MS

Crop Growth and Development and Environment Species Variability

- ✓ Effects of multiple environmental factors on crop growth and phenology across many important crop species.
- ✓ Can we use environmental productivity index concept across the species to quantify the responses and to develop functional relationships?

Crop Growth and Development and Environment Major Crops – Global Statistics - 2008

| Crop | Area, Million ha | Production, Million Mt. | Productivity, Mt. ha ⁻¹ |
|------------|------------------|-------------------------|------------------------------------|
| Wheat | 224 | 690 | 3.0 |
| Maize/Corn | 161 | 823 | 5.1 |
| Rice | 159 | 685 | 4.3 |
| Soybeans | 97 | 231 | 2.4 |
| Barley | 57 | 158 | 2.8 |
| Sorghum | 45 | 65 | 1.5 |
| Millets | 37 | 36 | 0.96 |
| Seedcotton | 31 | 66 | 2.1 |
| Rapeseed | 30 | 58 | 1.9 |
| Beans, dry | 28 | 20 | 0.73 |

Environmental and Cultural Factors Influencing Crop Phenology


- Atmospheric Carbon Dioxide (indirect)
- Solar Radiation (indirect)
- Photoperiod (direct on flowering, no effect on modern cotton cultivars)
- Temperature (direct)
- Water (indirect)
- Wind (indirect)
- Nutrients (N, P and K) (direct & indirect)
- Growth Regulators (PIX) (indirect)






Rice – Some Crop Statistics



- Provides the dietary needs of 1.6 billion with another 400 million rely on rice for one-quarter or one-half of their diet.
- **2004 stats are:** Area = 151 Million ha, production = 606 Million Mt, and average yield = 4.02 t ha⁻¹.
- 53% Irrigated flooded-paddy
- 27% Rainfed lowland
- 12% Rainfed upland
- 8% Deep-water

Rice Production Statistics











Corn or Maize, *Zea mays* L.

Corn is the 3rd most important food crops globally in terms of energy and protein (FAO, 2004).

Area: 147 Million ha
 Total production: 721 Million Mt
 Average yield: 4.93 t ha⁻¹



Phenology – Species Variability Sorghum, *Sorghum bicolor* (L.) Moench

Sorghum is the 4th most important food crops globally in terms of energy and protein (FAO, 2004).

2004 stats are:

Area: 43.1 Million ha
Total production: 57.8 Million Mt
Average yield: 1.3 t ha⁻¹



United States = 2.64 Mha, 4.4 t ha⁻¹, 11 MMt
India = 9.4 Mha, 0.8 t ha⁻¹, 7.53 MMt
Nigeria = 7.1 Mha, 1.13 t ha⁻¹, 8.03 MMt
China = 0.57 Mha, 4.1 t ha⁻¹, 2.34 MMt
Mexico = 1.91 Mha, 3.35 t ha⁻¹, 6.4 MMt
Sudan = 6.0 Mha, 4.33 t ha⁻¹, 2.6 MMt

Sorghum – Cultivar Differences

Phenology – Panicle Initiation.

- Notice the base temperature which is almost similar to corn.
- The rate of development can be described by a bilinear model.




Fig. 1. Relationship between the rate of progress from planting towards PI and mean temperature in sorghum genotypes IS32520 and IS11758 grown in controlled temperature glasshouses under short-days. Fitted lines where $T_1 < T \leq T_2$ $y = -0.021 - 0.0020x$; $T_2 < T \leq T_3$ $y = 0.114 - 0.0031x$ and where $T_1 = T_2 = 10^\circ\text{C}$, $T_2 = T_3 = 25^\circ\text{C}$.

Sorghum – Phenology – Cultivar Differences

Phenology – Leaf development.

- Notice the base temperature which is almost similar to corn.
 - The rate of development can be described by a bilinear models.
- Also, notice the genotypic variability.






Fig. 4. Relationship between rate of leaf-tip appearance and mean temperature in sorghum genotypes RS610 and IS25557 grown in controlled temperature glasshouses under short-days. Fitted lines where $T_1 < T \leq T_2$ $y = -0.148 - 0.018x$; and where $T_2 < T \leq T_3$ $y = 0.660 - 0.01215x$ and where $T_1 = T_2 = 10^\circ\text{C}$, $T_2 = T_3 = 25^\circ\text{C}$.


Sorghum – Vegetative Growth and Development



Sorghum – Vegetative and Reproductive Growth and Development



Sorghum – Reproductive Growth and Development







Wheat – Some Crop Statistics

- Provides 20% of the energy and 25% of the protein requirements of over 6 billion population.
- **2004 stats are:**
 Area = 217 Million ha
 Production = 633 Million Mt
 Average yield = 2.84 t ha⁻¹.


Wheat – Production Trends



Wheat – Production Trends





Wheat – Production Trends



Wheat and Phenology




Temperature Effects on Flowering to maturity



Soybean Production Statistics



Soybean the most important protein and oilseed crop globally (FAO, 2004).

Area: 91.4 Million ha
 Total production: 204.4 Million Mt
 Average yield: 2.23 t ha⁻¹




Temperature and CO₂ Effects on Soybean Developmental Rates

| CO ₂ , ppm | Temp. °C | Plastochron interval, d leaf ⁻¹ | Final node number, no. plant ⁻¹ |
|-----------------------|----------|--|--|
| 300 | 26/19 | 4.2 | 10.3 |
| | 31/24 | 3.3 | 11.5 |
| | 36/29 | 3.2 | 12.0 |
| 600 | 26/19 | 3.9 | 11.2 |
| | 31/24 | 2.7 | 11.4 |
| | 36/29 | 2.6 | 12.1 |




Temperature and Soybean Developmental Rates




Temperature and Soybean Developmental Rates


Flowering to seed maturity (R1 to R8)



Temperature and Soybean Growth Rates




Rangeland grass – Big bluestem (*Andropogon gerardii*)




- C4 photosynthetic pathway
- Clump forming perennial grass
- Grows 3 to 6 feet tall but occasionally up to 9 feet.
- Lower stems are a purplish or bluish color
- Leaves are 1/2 inch wide and up to 20 inches long.
- Arrangement of the flowers in three dense elongate clusters is the reason for the common name of turkey-foot grass.
- Grows best in moist well drained soil in full sun and is a major component of the tallgrass prairie.

Rangeland grass – Big Bluestem, *A. girardii* Reproductive Response – Panicle initiation




Rangeland grass – Big Bluestem, *A. girardii*

Growth and Developmental Responses






Rangeland grass – Big Bluestem, *A. girardii*



Rangeland grass – Big Bluestem, *A. girardii*

Seed number and weight response

Total biomass response



Chickpea is a cool-season crop grown substantially in South and West Asia, the Mediterranean region, and South and Central America.




Chickpea - *Cicer arietinum* L.

Production and distribution

- Chickpea is a cool season food legume crop grown on > 10 million ha in 45 countries of the world.
- Chickpea is either the first or the second most important, rainfed, cool season food legume, grown mainly by small farmers in the semi-arid tropics (SAT) and West Asia and North Africa (WANA) regions.
- The crop is also grown in southern and eastern Africa (particularly important in Ethiopia), Europe, the Americas and, more recently, Australia.
- World production is 7 million tones.
- International trade in chickpeas has increased over the years.


Phenology - Species Variability Temperature - Sowing to 50% Emergence Chickpea Cultivars




Roberts et al., 1980. Expt. Agric. 16:343-360.



Phenology - Species Variability Temperature - Sowing to First Flower Chickpea Cultivars


Roberts et al., 1980. Expt. Agric. 16:343-360.







Groundnut or Peanut
(Arachis hypogaea L.)

Groundnut, an important cash crop, is an annual legume. Its seeds are a rich source of edible oil (43-55%) and protein (25-28%). About two thirds of world production is crushed for oil and the remaining one third is consumed as food. Its cake is used as feed or for making other food products and haulms provide quality fodder.



Groundnut or Peanut
(Arachis hypogaea L.)










The semi-arid tropics of Asia and Africa (WC = western and central Africa; SEA = southern and eastern Africa (Source: <http://www.cgjar.org/icrisat/>))

Distribution:
 Groundnut originated in the southern Bolivia/north west Argentina region in south America and is presently cultivated in 108 countries of the world.


Asia with 63.4% area produces 71.7% of world groundnut production followed by Africa with 31.3% area and 18.6% production, and North-Central America with 3.7% area and 7.5% production.

Important groundnut producing countries are China, India, Indonesia, Myanmar, Thailand, and Vietnam in Asia; Nigeria, Senegal, Sudan, Zaire, Chad, Uganda, Cote d'Ivory, Mali, Burkina Faso, Guinea, Mozambique, and Cameroon in Africa; Argentina and Brazil in South America and USA and Mexico in North America.





Phenology and Growth - Species Variability Cardinal Temperatures - Major Crops



Sanchez et al. (2014), Global Change Biology, 20:408-417

Phenology and Growth - Species Variability Cardinal Temperatures - Major Crops


Table 1. Summary of mean (SE) and leaf initiation (T_{min}) and leaf senescence (T_{max}) temperatures (T_{min} optimum (T_{opt}) and maximum (T_{max}) temperature for different phenology and development phases in terms of number of calendar months.

| Phenology | Mean Temperature (SE) (°C) | T _{min} (°C) | T _{opt} (°C) | T _{max} (°C) |
|-----------------|----------------------------|-----------------------|-----------------------|-----------------------|
| Leaf initiation | | | | |
| Rice | 18.0 (0.5) | 10 | 25 | 30 |
| Wheat | 16.0 (0.2) | 8 | 22 | 28 |
| Maize | 21.0 (0.3) | 12 | 28 | 35 |
| Shoot growth | | | | |
| Rice | 24.0 (0.3) | 15 | 30 | 35 |
| Wheat | 22.0 (0.2) | 12 | 28 | 32 |
| Maize | 28.0 (0.4) | 20 | 35 | 40 |
| Root growth | | | | |
| Rice | 18.0 (0.3) | 10 | 25 | 30 |
| Wheat | 16.0 (0.2) | 8 | 22 | 28 |
| Maize | 21.0 (0.3) | 12 | 28 | 35 |
| Leaf senescence | | | | |
| Rice | 30.0 (0.4) | 25 | 35 | 40 |
| Wheat | 28.0 (0.3) | 22 | 32 | 38 |
| Maize | 35.0 (0.5) | 30 | 40 | 45 |
| T germination | | | | |
| Rice | 10.0 (0.2) | 5 | 15 | 20 |
| Wheat | 8.0 (0.1) | 3 | 12 | 18 |
| Maize | 12.0 (0.3) | 7 | 18 | 25 |
| T optimum | | | | |
| Rice | 25.0 (0.3) | 15 | 30 | 35 |
| Wheat | 22.0 (0.2) | 12 | 28 | 32 |
| Maize | 28.0 (0.4) | 20 | 35 | 40 |
| T maximum | | | | |
| Rice | 30.0 (0.4) | 25 | 35 | 40 |
| Wheat | 28.0 (0.3) | 22 | 32 | 38 |
| Maize | 35.0 (0.5) | 30 | 40 | 45 |


Sanchez et al. (2014), Global Change Biology, 20:408-417

Phenology and Growth - Species Variability Temperatures Response Functions - Major Crops

Warm-season Turf Grasses



Cool-season Turf Grasses



Phenology – Species Variability Horticultural Crops



Phenology – Species Variability




Fig. 10.3. (a) Relationship between temperature and the rate and percentage of germination of onion seeds on moist paper. Rates are reciprocals of the number of days for 50% of viable seeds to germinate (data of Harrington, 1962). (b) Relationship between temperature and rate of cotyledon elongation before hook formation for newly germinated onion seedlings cv. White Lisbon (Wheeler and Ellis, 1991). (c) Relationship between temperature and relative rate of cotyledon elongation after hook formation for the same seedlings as in (b) (Brewster, 1997).

Phenology – Species Variability – Crops and weeds

| Species | T _{min} or T _{base} , °C | GDD per leaf tip |
|---------------------|--|------------------|
| Maize | 8 | 39 |
| Sorghum | 8 | 48 |
| Pearl millet | 12 | 26 |
| Wheat | 0 | 99 |
| Barley | 1 | 75 |
| Rice | 5 | 90 |
| Soybean | 7 | 54 |
| Sunflower | 9 | 29 |
| Cowpeas | 16 | 30 |
| Sugar beet | 2 | 30 |
| Velvet leaf | 8 | 24 |
| Pigweed | 10 | 12 |
| Banana | 8 | 196 |

Phenology – Species Variability

Phenology and Seed Germination:

- The minimum or the base temperature, heat sum (s) or the growing degree days (GDD) from that base temperature for a number of horticultural crops.

Table 10.2. Minimum germination temperature (T_{min}) and heat sum (S) in degree-days for seedling emergence, and the applicable temperature (T) range for germination of various vegetables. Crops are ranked within groups by heat sum (S) in degree-days. (from Taylor, 1987.)

| Group | Crop | Genus and species | T_{min} (°C) | S (degree-days) | T (°C) |
|------------------------------------|------------------|--|--------------------|-----------------|--------|
| Leaf vegetables and brassica crops | Broccoli | <i>Brassica oleracea</i> | 11.0 | 48 | 15-25 |
| | Cress | <i>Lepidium sativum</i> | 1.0 | 64 | 3-17 |
| | Lettuce | <i>Lactuca sativa</i> | 3.5 | 71 | 6-21 |
| | Witloof, Chicony | <i>Cichorium endivia</i> | 5.3 | 85 | 9-25 |
| | Endive | <i>Cichorium endivia</i> | 2.2 | 93 | 3-17 |
| | Spiny cabbage | <i>B. oleracea</i> var. <i>cauliflora</i> | 1.9 | 95 | 3-17 |
| | Turnip | <i>B. campestris</i> var. <i>rapa</i> | 1.4 | 97 | 3-17 |
| | Broccoli, kale | <i>B. oleracea</i> var. <i>capitata</i> | 1.2 | 103 | 3-17 |
| | Red cabbage | <i>B. oleracea</i> var. <i>purpuracea</i> | 1.3 | 104 | 3-17 |
| | White cabbage | <i>B. oleracea</i> var. <i>capitata</i> | 1.0 | 106 | 3-17 |
| | Broccoli sprouts | <i>B. oleracea</i> var. <i>gemmifera</i> | 1.1 | 108 | 3-17 |
| | Spinach | <i>Spinacea oleracea</i> | 0.1 | 111 | 3-17 |
| Fruit vegetables | Cauliflower | <i>B. oleracea</i> var. <i>botrytis</i> | 1.3 | 112 | 3-17 |
| | Corn salad | <i>Valerianella olitoria</i> | 0.0 | 141 | 3-17 |
| | Leek | <i>Allium porum</i> | 1.7 | 222 | 3-17 |
| | Celery | <i>Apium graveolens</i> | 4.6 | 237 | 9-17 |
| | Parsley | <i>Petroselinum crispum</i> | 0.0 | 240 | 3-17 |
| | Tomato | <i>Lycopersicon esculentum</i> | 8.7 | 88 | 13-25 |
| | Aubergine | <i>Solanum melongena</i> | 12.1 | 93 | 15-25 |
| | Chickpea | <i>Cicerum sativum</i> | 12.1 | 108 | 15-25 |
| | Melon | <i>Cucumis melo</i> | 12.2 | 108 | 15-25 |
| | Sweet pepper | <i>Capiscum annuum</i> | 10.9 | 122 | 15-25 |
| | Carrot | <i>Daucus carota</i> | 1.3 | 170 | 3-17 |
| | Leguminous crops | Onion | <i>Allium cepa</i> | 1.4 | 219 |
| French sugar pea | | <i>Pis. sativum</i> var. <i>sacharatum</i> | 1.6 | 96 | 3-17 |
| Bean (French) | | <i>Phaseolus vulgaris</i> | 7.7 | 139 | 13-25 |
| Broad bean | | <i>Vicia faba</i> | 0.4 | 148 | 3-17 |
| Radish | | <i>Raphanus sativus</i> | 1.2 | 75 | 3-17 |
| Scaroletera | | <i>Scorzonera hispanica</i> | 2.0 | 90 | 3-17 |
| Beet | | <i>Beta vulgaris</i> | 2.1 | 119 | 3-17 |
| Carrot | | <i>Daucus carota</i> | 1.3 | 170 | 3-17 |
| Onion | | <i>Allium cepa</i> | 1.4 | 219 | 3-17 |

EPI Concept and Plant Growth and Development

One way to quantify the effects of environmental factors on plant growth and development is to use the EPI concept similar to the one that we used in cotton as model crop.

EPI-phenology = Temperature (potential) * Nutrient Index (C, N, P, K) * Water index * PPF Index * PGR Index etc.,

EPI-growth = Temperature (potential) * Nutrient Index (C, N, P, K)* Water index * PPF Index * PGR Index etc.,

Once the potential is defined and quantified, then we can use EPI concept to decrease that potential to account for the effects of multiple environmental factors on given process such growth or development of any plant/crop species as in cotton crop.

EPI Concept and Plant Growth and Development

- Environmental productivity index concept, if applied, works across species and locations.
- EPI also allows one to interpret and to understand stresses in the field situations.
- If we know the factor that is limiting most at any point of time during the growing season, then, we can make appropriate management decisions to correct that limitation.
- EPI concept is the way to quantify the effects of multiple environmental factors on plant growth and development (photosynthesis, phenology, and growth) and thus productivity of any species or crop.