

Environmental Plant Physiology Summary


K. Raja Reddy
Mississippi State University
Mississippi State, MS

Environmental Plant Physiology Objectives


- The objectives of this course were to learn plant responses to abiotic stresses, particularly plant growth and development, and to learn modeling methodologies on how to integrate those plant processes under multiple stress conditions.
- At the end, the students were expected to:
 - ✓ understand individual as well as interactive abiotic stress effects on photosynthesis, respiration, growth, development and finally yield.
 - ✓ understand on how to develop methodologies to integrate multiple stress factor effects on various plant/canopy processes.

Trends That Shape Our Future


Trends, Signs and Signatures from the Earth Past, Present and Future World Population



Trends, Signs and Signatures from the Earth Present and Future World Population Trends




Trends, Signs and Signatures from the Earth Global Major Foods – Per Capita Consumption




Trends, Signs and Signatures from the Earth Global Major Foods – Meat and Poultry Production


Meat and Poultry Production



Meat and Poultry Production
Relative Trends




Trends, Signs and Signatures from the Earth Maize - Production and Yield – Selected Counties




Trends, Signs and Signatures from the Earth Management Practices

Crop rotations



Fertility management




Trends, Signs and Signatures from the Earth Cropland area, Irrigation and Salinization


	Year 2000		
	Cropland area	Irrigated area	Salinized area
	----- Mha -----		
China	124.0	54.4 (22%)	7-8 (14%)
India	161.8	54.8 (31%)	10-30 (50%)
USA	177.0	22.4 (13%)	4.5-6 (15%)
USSR	204.1	19.9 (2%)	2.5-4.5 (21%)
World	1364.2	271.7 (21%)	62-82 (37%)

Percent change since 1985

Trends, Signs and Signatures from the Earth Population, cereal yield, arable and irrigated area, N use



Trends, Signs and Signatures from the Earth Yield Trends of Major Crops – Past and Future



- Yield has to be doubled by 2050 to meet the demands of rising population with higher earning capacity, diet shifts, and increasing biofuels consumption.
- Based on past trends, projected rate of yield increase is about 1.2% per year, but we need about 2.4% per year to double the current yields for major cereal crops including rice.

Ray et al., 2013, PLOSOne

Feeding 10 Billion Mouths

We must develop the capacity to feed over 10 billion people within in the next 35 to 85 years.


- The average world's current cereal yield is about **3.89 tons per ha** for about **7.2 billion people** (year 2014).
- We need about 5.24 tons per ha for 9.7 billion (2050; 35 % more than the current), and 5.94 tons per ha for 11 billion (2100; 53% more than the current).

Routes to Greater Food Production

- Increase in the area of land under cultivation.
- Increase in the number of crops per hectare per year (mostly practiced in tropics, requires access to irrigation, high input use, short season cultivars, and others such as labor, pest and disease control may be a problem).
- Displacement of lower yielding crops by higher yielding ones (done since the dawn of domestication).
- Efficiency of crop production in terms of:
 - Per unit of land area (yield per ha)
 - Per unit of time
 - Per unit of inputs such as fertilizers, water and labor etc.

Here comes the greatest challenge of our time,
The Global Climate Change

Trends, Signs and Signatures from the Earth
In a production environment, no two seasons are equal



Environmental Stresses and Plant Growing Conditions

Environmental and Cultural Factors Limiting Potential Yields

- Atmospheric carbon dioxide
- Solar radiation
- Temperature (extremes)
- Water (irrigation and rainfall)
- Wind
- Nutrients (N, P, K, and other nutrients)
- Others, Ultra-violet radiation, ozone etc.,
- Growth regulators (such as PIX)

Area of Total World Land Surface Subject to Environmental Limitations of Various Types

Limitation	Area of world soil subject to limitation (%)
Drought	27.9
Shallow soil	24.2
Mineral excess or deficiency	22.5
Flooding	12.2
Miscellaneous	3.1
None	10.1
Total	100
Temperature	14.8 (over laps with other stresses)

Environmental Plant Physiology

Chapter 1:

- Atmospheric carbon dioxide
- Solar radiation
- Temperature (Including extremes)
- Water (rainfall, flooding, and irrigation)
- Wind
- Nutrients
- Other factors such as ozone
- Plant growth regulators
- The facilities and tools

Environmental Plant Physiology

Chapter 2:

Photosynthesis and the environment

- The Environmental productivity index (EPI) concept.
- The photosynthesis - Species variability.
- Photosynthesis and aging process.
- Respiration.

Environmental Plant Physiology

Chapter 3:

Crop growth and development

- Phenology
- Growth of various organs and whole plants.
- The concept of environmental productivity index in quantifying crop growth and development in response to the environment.

Environmental Plant Physiology

Chapter 4:

Scaling of processes from leaves to whole plant, canopies or ecosystems.

Chapter 5:

Special topics include:




- Plant growth regulators – PIX.
- Remote sensing and environmental plant physiology.

Environmental limiting crop growth, development and yield

- Atmospheric Carbon Dioxide
- Solar Radiation
- Temperature
- Water (indirect)
- Wind
- Nutrients (N, P, K)
- Ozone, UV-B etc.,
- Growth Regulators

Environmental limiting crop growth, development and yield

- Atmospheric Carbon Dioxide
- Solar Radiation
- Temperature
- Water (indirect)
- Wind
- Nutrients (N, P, K)
- Ozone, UV-B etc.,
- Growth Regulators



Environmental limiting crop growth, development and yield





- Atmospheric Carbon Dioxide
- Solar Radiation
- Temperature
- Water (indirect)
- Wind
- Nutrients (N, P, K)
- Ozone, UV-B etc.,
- Growth Regulators

Radiation and Plant Life


Table 1.6. Effect of radiation on plant life. (Ross 1981)



Spectral region	Wavelength (nm)	Percent of solar radiant energy	Photo-synthetic	Effects of radiation		
				Photo-morpho-genetic	Photo-destruc-tive	Thermal
Ultraviolet	290-380	0-4	Insignifi-cant	Slight	Signifi-cant	Insignifi-cant
Photosynthetically active range (PhAR)	380-710	21-46*	Signifi-cant	Signifi-cant	Slight	Significant
Infrared	750-4000	50-79*	Insignifi-cant	Signifi-cant	Insignifi-cant	Significant
Longwave radiation	4000-100000		Insignifi-cant	Insignifi-cant	Insignifi-cant	Significant

* Depending on position of sun and degree of cloud cover.



- ### Effects of Multiple Environmental Factors on Crop Growth and Developmental Aspects
- Introduced Environmental Productivity Index (EPI) concept.
 - Photosynthesis
 - Crop Phenology or Development
 - Crop Growth
 - Reproductive Biology





Radiation Totals for the 1992 Growing season

Mississippi State – North Farm

Variable	Amount, MJ
Total Incoming Radiation	2842
Intercepted Radiation	1551
Percent Intercepted	55

Photosynthesis – EPI Concept

Accounting for Individual factors

Variable	Amount, g CO ₂ m ⁻² season ⁻¹
Incoming R	19644
Intercepted R	11441 (100%)
Int. R * UV-B	10448 (9%)
Int. R * T	10139 (11%)
Int. R * W	9783 (14%)
Int. R * N	8986 (21%)
Int. R * K	10841 (5%)

Photosynthesis – EPI Concept





Accounting for Multiple Factors

Variable	Amount, g CO ₂ m ⁻² season ⁻¹
Incoming R	19644
Intercepted R	11441 (100%)
Int. R * UV-B	10230 (9%)
Int. R * UV-B * T	9153 (20%)
Int. R * UV-B * T * W	7551 (34%)
Int. R * UV-B * T * W * N	6292 (55%)
Int. R * UV-B * T * W * K	4576 (60%)

Actual amount

Environmental limiting crop growth, development and yield

- Atmospheric Carbon Dioxide
- Solar Radiation
- Temperature
- Water (indirect)
- Wind
- Nutrients (N, P, K)
- Ozone, UV-B etc.,
- Growth Regulators



Environment Factors

Temperature:

- **Strongly Affects:**
 - Phenology
 - Vegetative growth, LAI, LAD
 - Fruit Growth and Retention
 - Respiration
 - Water-loss and Water-Use
- **Moderately Affects:**
 - Photosynthesis on a canopy basis

High-temperature Injury


Heat-blasted Cotton Flowers and Squares – Arizona

Heat-blasted Cotton Squares
California's San Joaquin Valley

Figure 7. Heat-blasted squares in California's San Joaquin Valley. (Photo: R. Vargas)

Environmental limiting crop growth, development and yield

- Atmospheric Carbon Dioxide
- Solar Radiation
- Temperature
- Water (indirect)
- Wind
- Nutrients (N, P, K)
- Ozone, UV-B etc.,
- Growth Regulators



Water

Water plays essential roles in plants as a:


- Constituent
- Solvent
- Reactant in various chemical processes
- Maintenance of turgidity

The physiological importance of water is reflected in its ecological importance.


The distribution plants over the earth's surface is controlled by the availability of the water (amount and seasonal distribution of precipitation) where ever temperature permits growth.

Figure 10

Relative Precipitation Changes (%/°C) Mean of 15 Models



Role of water in cotton



Environment Factors

Water Deficits:

- Strongly affects:
 - Vegetative growth, LAI, LAD
 - Fruit Growth and Retention
 - Water-loss and Water-Use
 - Photosynthesis
- Moderately affects certain phenological events:
 - Phenology (leaf development)


Environment – Wind and Hail




Environmental limiting crop growth, development and yield

- Atmospheric Carbon Dioxide
- Solar Radiation
- Temperature
- Water (indirect)
- Wind
- Nutrients (N, P, K)
- Ozone, UV-B etc.,
- Growth Regulators

Photosynthesis - Variability Among Species Response to Leaf Nitrogen



Environment - Nitrogen



Environment - Nitrogen

Parameter	Percent Reduction from the Optimum (2.5 g N/m ² or 4.5%)			
Leaf N, g m ⁻²	Photosynthesis	Stem growth	Leaf growth	Leaf Development
2.5	100	100	100	100
2.0	12	14	18	12
1.5	53	60	>99	68
1.2	76	--	--	--

Environment - Nitrogen




Figure 2 The effect of N on number of main-stem nodes, boll number per plant, leaf area, and boll weight (Jackson and Gerik, 1990).


Environment Factors

Fertilizers Deficits - Potassium:

- **Strongly Affects:**
 - Leaf growth, LAI, LAD
 - Fruit Retention
- **Moderately Affects:**
 - Photosynthesis
 - Stem growth

Environment - Nutrients

Potassium - Cotton Growth and Development Environmental Productivity Indices



Environment Factors

Ultraviolet-B Radiation:

- **Strongly Affects:**
 - Photosynthesis
 - Stem growth
- **Moderately Affects:**
 - Leaf growth
 - Leaf aging
- **No Effects:**
 - Phenology

Environmental limiting crop growth, development and yield

- Atmospheric Carbon Dioxide
- Solar Radiation
- Temperature
- Water (indirect)
- Wind
- Nutrients (N, P, K)
- Ozone, UV-B etc.,
- Growth Regulators

Solar Radiation and Plant Life

For plants, radiation is:


- A source of energy (photoenergetic effect).
- Stimulus for development (photocybernetic effect).
- Stress factor (photodestructive effect).

Effects of Radiation on Plant Life

Spectral Region	Wavelength nm	%	Photo-synthe-sis	Effects		
				Photo-morpho-genetic	Photo-destro-ctive	Thermal
Ultraviolet	290-380	0-4	IS	Slight	S	IS
PAR	380-710	21-46	S	S	Slight	S
Infrared	750-4000	50-79	IS	S	IS	S
Longwave	4000-100000		IS	IS	IS	S

IS = Insignificant, S = Significant


Ultraviolet Radiation




- **UVC: <280), UVB: 280-320, and UVA: 320-400.**
- UVA is not absorbed by ozone.
- UVB is mostly absorbed by ozone, although some reaches the Earth.
- UVC is completely absorbed by ozone and normal oxygen.

Why are we concerned with UV now?


The Ozone Depletion Process




Satellite Based Ozone levels - 2001




Satellite Based UV Radiation levels - 2001




UV-B Radiation – Growth EPI Factors for various growth Processes



UV-B Radiation – Phenology EPI Factors for various Developmental Processes



UV-B Radiation – Cotton Growth EPI Factors for various growth Processes



Environmental limiting crop growth, development and yield




- Atmospheric Carbon Dioxide
- Solar Radiation
- Temperature
- Water (indirect)
- Wind
- Nutrients (N, P, K)
- Ozone, UV-B etc.,
- Growth Regulators

Environment Factors


Growth Regulators - Mepiquat Chloride (PIX):

- **Moderately Affects:**
-- Leaf, stem and branch growth and LAI
- **Slightly Affects:**
-- Photosynthesis


Mepiquat Chloride (PIX) - Growth EPI Factors



Flow Diagram of GOSSYM Cotton Crop Simulation Model



Technology Fusion and Delivery System



Environmental Plant Physiology and Crop Modeling

- Modeling forces the organization of known information and concepts.
- Although we may not know enough to develop a comprehensive model that includes all aspects of the farm or crop production system, modeling some meaningful portions of the system provides clarity.
- For a model to correctly predict plant responses to physical conditions, the concepts and the response functions must be appropriately assembled.

Environmental Plant Physiology, Crop Modeling, and Technology Integration for Decision Support System

- Critical environment-genotype relations should be incorporated into the model.
- When a crop model is based on appropriate concepts and processes it will have the predictive capability in new environments, and can be used either alone or with other emerging newer technologies to disseminate useful production information.
- Also, crop models should be integrated with other related technologies for technology integration and delivery.

Environmental Plant Physiology Summary and Conclusions

- To study the effects of environmental factors on growth, development and other processes, we need:
 - ✓ Controlled environmental facilities with realistic environmental conditions including solar radiation.
 - ✓ Breakdown of whole systems into sub-systems and study the influence of environmental factors on those subsystems.
 - ✓ Develop some concepts such as EPI to quantify the effects of multiple environmental factors on subsystems.
 - ✓ Integrate sub-systems into coherent whole plant/field/ecosystem system-level models/tools.

Environmental Plant Physiology Summary and Conclusions

- Validated/integrated system simulation models will be useful:
 - ✓ To test hypothesis.
 - ✓ To understand multiple environmental effects or interactions.
 - ✓ Can be used for resources management at the field-level.
 - ✓ Can be used for resource management to assist policy decisions.
 - ✓ Can be used as an educational tool to understand the effects of environment/management effects on crop functioning.
 - ✓ Can be used for impact assessment of climate change on crop production systems across regions and nations.

“You cannot build peace on empty stomachs.”

John Boyd Orr
Nobel Peace Laureate
First FAO Director General



“You can’t eat the potential yield, but need to raise the actual by combating the stresses”

Norman E. Borlaug
Nobel Peace Laureate

