


Crop Growth

High-temperature Injury to Reproductive Parts

K. Raja Reddy
Mississippi State University
Mississippi State, MS

Plant Responses to Extreme Temperatures



**Plants lack locomotion
They should either adapt or tolerate stress**

Plant Responses to Extreme Temperatures

- Few plant species survive a steady high temperatures above 45 °C
 - ✓ Actively growing tissues can rarely survive over 45°C
 - ✓ However, non-growing cells or organs (Pollen and seed) can survive much higher temperatures.
 - some pollen up to 70 °C
 - some seed up to 120 °C.
- Heat stress is also a major problem in greenhouses, where low air speed and high humidity decreases leaf cooling and thus affecting leaf/canopy temperatures.

Plant Responses to Extreme Temperatures

- Plants do adapt to high temperature:
 - ✓ Reflective leaf hairs and waxes
 - ✓ Leaf rolling, and vertical leaf orientation
 - ✓ Small leaves and dissected (okra) leaf morphology
 - ✓ Synthesis of heat-shock proteins (HSPs)
 - ⇒ Help cells withstand heat stress
 - ⇒ However, the functions of all HSPs are not yet fully known, but many act as molecular chaperons, help stabilize and fold proteins, assist in polypeptide transport across membranes, protect enzymes, etc.


Plant Critical Processes at Extreme Temperatures

- Photosynthesis and respiration, and conductivity will be affected by high temperatures.
- However, photosynthesis declines faster than respiration and conductivity at high temperatures.
- The point when the amount of CO₂ fixed equals to the amount of CO₂ released by respiration is called temperature compensation point. At this point and beyond, the carbon is not replaced, and carbohydrate reserves will be used for cellular functions.
- Therefore, the imbalance between photosynthesis and respiration causes deleterious effects at high temperatures.


Plant Critical Processes at Extreme Temperatures

- The question is how do plant groups respond to high temperatures?
- Enhanced temperatures are more detrimental in C_3 plants than in C_4 or CAM plants because of rates of both dark and photorespiration are increased more in C_3 plants.
- What happens to C_3 plants under elevated CO_2 conditions?


Climatic Zones and Temperature Conditions




Long-Term Average Temperatures




Long-term Average Temperatures for Four US Cotton Producing Areas



Hourly Temperatures for July 1995 Mississippi State, MS




Temperature Conditions - Diurnal Trends Mississippi State, MS - 1995



Diurnal temperature data recorded in June 1989 at Fatehpur, Rajasthan, India, (Latitude 27°C 37'N).

- 5 cm depth of soil (▲)
- 0.5 cm depth of soil (●)
- 150 cm above the soil surface (■)




Howarth, 1991

Climate Change and Crop Production

- Past changes in greenhouse gases has resulted about 0.6 °C increase in global average temperature during the last century.
- If current and future rates of changes in greenhouse gases and other land-use changes continue, then, these changes will exacerbate the natural climate changes and may result in:
 - 2 to 6 °C warmer temperatures
 - More frequent episodes of extreme events (heat, cold, drought, excessive rainfall resulting in floods, severe hurricanes, etc.).

Second green revolution to overcome environmental stresses



High Temperature Effects on Cotton Fruit Production and Retention

Pima Cotton Responses to Temperatures

The next 3 video clips show cotton responses to optimum (30/22°C, day/night), higher (35/27°C) and super-optimum (40/32°C) temperatures.

Notice that the plants grown in optimum temperatures are producing both vegetative and reproductive structures continuously and there is no abscission of squares or fruiting structures. Plants grown in 35/27°C are producing luxuriant vegetative growth, but some of the squares are being abscised due to excessive heat. If plants are grown in 40/32°C, the vegetative growth is reduced to certain extent compared to plants grown in other temperatures, but there is a complete reproductive failure (no flower-bud initiation and even fruiting branch production) due to excessive heat.

Optimum Temperature No Injury to Reproductive Parts




Higher Temperature Injury
 Partial Injury to Reproductive Parts




Super High Temperature Injury
 Total Reproductive Failure, Including
 Fruiting Branch Production



High Temperature Effects on Cotton – Upland Cotton



High Temperature Effects on Cotton – Pima Cotton




Environment - Crop Growth – High Temperature Injury to Reproductive Parts

Table 1. Effect of temperature on cotton growth, cv. Stoneville 825, harvested 49 days after initiation of temperature treatments. treatments are imposed at first flower. Standard error of the mean values are shown.

	Day/Night Temperature, °C				
	20/12	25/15	30/20	35/25	40/30
	Grams per Plant				
Total Wt.	242	320	330	293	225
% of Optimum	73	97	100	88	68
Bolls	17	63	143	17	0.8
% of Optimum	12	44	100	12	0.6

Environment - Crop Growth – High Temperatures Injury to Reproductive Parts




Projected Temperatures and Cotton Development


Treatment	Days to the Event		
	Square	Flower	Open Boll
1995 minus 2°C	33	65	144
1995 plus 0°C	26	51	101
1995 plus 2°C	24	48	94
1995 plus 5°C	21	42	77
1995 plus 7°C	19	39	No Fruit

No significant differences were observed between CO₂ levels


High Temperature Injury Temperature and CO₂ Interactions




High Temperature Injury Temperature and CO₂ Interactions – Cotton




High Temperature Injury Temperature and CO₂ Interactions – Cotton





High Temperature Injury Temperature and CO₂ Interactions – Cotton



High Temperature Injury Temperature and CO₂ Interactions – Cotton





High Temperature Effects on Cotton

The high temperature injury in cotton to reproductive growth and development is not fully understood so far.

High temperature causes some heat-sensitive cultivars/species (Pima cotton) to be vegetative (total reproductive failure and the reproductive induction process is sensitive). Not much is known why plants stay vegetative at those high temperature conditions.

Once the flower-buds (squares) are formed, exposure to extremely high temperatures (35/27°C) will result in abscission of squares.

High Temperature Effects on Cotton

Nutrient starvation is not the factor that causes that square abortion because plants grown in elevated or twice ambient CO₂ and under optimum nutrient conditions also drop those squares, and the nutrient demand for squares is minimal.

The evidence suggest that the 2 weeks prior to and 1 week post flower is the most sensitive stage in cotton.

Systematic evaluation is needed to quantify the effects of high temperature on both the male (anther, pollen growth and development) and female (ovule growth and development).

High Temperature Effects on Cotton

Breeders need simple and quantitative methods to screen genotypic variability and to find or breed a genotype to a niche environment for optimum crop production.

Biotechnology may play a role in developing cultivars that are more heat-tolerant.


Heat-tolerance will be beneficial even in today's environment, and will be needed more in a warmer future climatic conditions.

High Temperature Target Sites in Plants

Morphological, physiological, metabolic, and reproductive changes in plants under heat stress.

(1) **Direct injury:** includes aggregation and denaturation of proteins as well as increased fluidity of membranes.


(2) **Indirect injury:** includes inactivation of enzymes in chloroplasts and mitochondria, inhibition of protein synthesis, enhanced protein degradation and loss of membrane integrity. All these alterations result in cell injury or even death within a few minutes, which ultimately leads to catastrophic collapse of cellular organization



Kaushal et al., *Cogent Food & Agriculture* (2016), 2: 1134380
<http://dx.doi.org/10.1080/23311932.2015.1134380>


High Temperature Target Sites in Plants

Both megagametophyte and microgametophyte are the main target sites of heat stress, with the former more heat sensitive.



Kaushal et al., *Cogent Food & Agriculture* (2016), 2: 1134380
<http://dx.doi.org/10.1080/23311932.2015.1134380>

High Temperature Effects on Reproductive and Grain-filling Stages




Kaushal et al., *Cogent Food & Agriculture* (2016), 2: 1134380
<http://dx.doi.org/10.1080/23311932.2015.1134380>

High Temperature Effects on Reproductive and Grain-filling Stages


Summary of HT injury to anther early development and effect of exogenous application of auxin.

The four differentiated layers of anther wall cells (epidermal, endothecium, middle layer, and tapetum cells) are sequentially degraded during pollen maturation. This degradation process appears to be controlled by programmed cell death (PCD) and results dehiscence of anther walls.




Higashitani, A. 2013. High-temperature injury and auxin biosynthesis in microsporogenesis. *Front. Plant Sci.* 4:47, 1-4.

DIVERSE RESPONSE OF PLANTS TO HEAT STRESS














Kaushal et al., *Cogent Food & Agriculture* (2016), 2: 1134380
<http://dx.doi.org/10.1080/23311932.2015.1134380>

High Temperature Injury Temperature and CO₂ – Rice Growth



Baker and Allen, 1993





Temperature Effects on Crop Yield Several Major Crops

Crop	Topt, °C	Tmax, °C	Yield at Topt, t/ha	Yield at 28 °C, t/ha	Yield at 32°C t/ha	% decrease (28 to 32 °C)
Rice	25	36	7.55	6.31	2.93	54
Soybean	28	39	3.41	3.41	3.06	10
Dry bean	22	32	2.87	1.39	0.00	100
Peanut	25	40	3.38	3.22	2.58	20
Grain sorghum	26	35	12.24	11.75	6.95	41

Allen et al., 2000


High Temperature Effects on Growth Stages of Major Crops

Table 3.4 High temperature effects on growth stages of major crops (from Acock and Acock, 1993)




Crop	Effects
Wheat	Temperature >30°C for >8 h, can reverse vernalization
Rice	Temperature >35°C for >1 h at anthesis causes spikelet sterility
Maize	Temperature >36°C causes pollen to lose viability
Soybean	Great ability to recover from stress. No especially critical period in its development
Potato	Temperature >20°C depresses tuber initiation and bulking
Cotton	Temperature >40°C for >6 h causes bolls to abort

- ### High Temperature Injury Conclusions – Temperature and CO₂ Interactions
- There are no beneficial effects of elevated CO₂ on reproductive processes.
 - There are no beneficial interaction of CO₂ on temperature effects on reproductive processes and yield.
 - Negative effects of elevated temperature on seed set, seed yield and harvest index were greater at elevated CO₂ (grain sorghum, dry bean and big blue stem).


High Temperature and Crop Productivity



Effects of Multiple Abiotic Factors



UV-B Radiation and Temperature Cotton Reproductive Growth and Development



UV-B Radiation and Temperature Cotton Reproductive Growth and Development



UV-B Radiation and Soybean Genotypes Reproductive Growth and Development


Treatments



Growing Conditions and Treatments:

Temperature (°C)	CO ₂ (ppm)	UV-B (kJ m ⁻² d ⁻¹)
30/22	360	0
		5
38/30	720	10
		15


UV-B and Temperature Soybean Reproductive Development – Sensitive Cultivar









Climate Change and Crop Productivity Conclusions – Temperature and CO₂ Interactions


- There are no beneficial effects of elevated CO₂ on reproductive processes in the crops investigated (cotton, soybean, rice, sorghum and beans).
- There are no beneficial interaction of temperature on UV-B effects on reproductive processes.
- High temperatures and higher UV-B aggravated the damaging effect on many reproductive processes.
- Elevated CO₂ did not ameliorate the damaging effects of either higher temperatures or elevated UV-B levels.

High Temperature and Crop Productivity



Genotypic Variability





- ### High Temperature Injury – Crop Reproduction Concluding Remarks
- The influence of stress factors on reproductive biology of crops/plants has not been well studied.
 - Better screening tools/methods are needed to assess the genotypic variability among crop species.
 - The current rate of climate change and climate variability and projected changes in climate are unprecedented, and plants may not cope with these rapid changes.
 - There is an urgent need to develop crop cultivars to a variety of stresses (high and low temperatures, water/drought stress, salt stress, UV-B radiation stress etc. either alone or in combination).