


Environmental productivity indices for crop growth and development: Cotton as an example

Photosynthesis



KRReddy@pss.MsState.edu
Department of Plant and Soil sciences

Mississippi State University
A Land-Grant Institution

Photosynthesis and Respiration and Environment

Goals and Learning Objectives:

- To understand the effects of multiple environmental factors on photosynthesis and respiration.
 - Photosynthesis and environment and Environmental Productivity Index (EPI) concept using cotton as an example crop.
 - Photosynthesis and environment and species variability and applicability of EPI concept.
 - Leaf and canopy aging and their relationship with photosynthesis.
 - Respiration and environment

Racing towards Enhancing Crop Photosynthesis

- The next advance in field crop productivity will likely need to come from improving crop resource use efficiencies (e.g. radiation, water, nutrients, etc.), which are linked with overall crop photosynthetic efficiency.
- For this, there is an emerging agenda focused on genetic manipulation of the biochemistry of photosynthesis process to enhance crop canopy photosynthesis, and thus productivity and yield.
- However, progress is limited by the lack of connection between biochemical/leaf-level photosynthetic manipulation and crop performance, which is influenced by genetics and plant growth and developmental processes and environmental effects.
- Crop models which can incorporate the interactions and integrate across scales of biochemical organization might be the tools needed to accelerate the process in photosynthetic enhancement.


Photosynthesis and Environment

You will learn:


- Effects of environmental factors on photosynthesis.
- How to quantify the effects of multiple environmental factors on photosynthesis.
- How to calculate potential photosynthesis under optimum conditions.
- Then, how to develop environmental productivity indices for various environmental factors to decrement the potential photosynthesis and to calculate actual photosynthesis.

Photosynthesis


- The process in which plants uses the energy from sunlight to combine carbon dioxide (CO₂) from the air with water to make carbohydrates plus oxygen.

$$6 \text{ CO}_2 + 6 \text{ H}_2\text{O} \xrightarrow[\text{Water, Nutrients}]{\text{Light, Plant}} \text{C}_6\text{H}_{12}\text{O}_6 + 6 \text{ O}_2$$


Environmental and cultural factors affecting Cotton growth and productivity




Global Atmospheric CO2 Concentrations Mauna Loa, HI and South Pole




Radiation Conditions - Seasonal Trends

Bakersfield, CA, Corpus Christi, TX and Stoneville, MS




Long-term Average Temperatures for Four US Cotton Producing Areas




Precipitation - Seasonal Trends

Bakersfield, CA and Florence, SC - 1991




Seasonal Trends - Midday Leaf Water Potential

Irrigated and rainfed cotton, MSU North Farm -1995




Seasonal Trends Solar and UV-B Radiation Mississippi State - 2001




Photosynthesis - Management Factors

➤ Management factors such as fertilizer application amounts and timings affect nutrient uptake and leaf nutrient status and thus photosynthesis (Leaf N, P, K etc..)


Cultural and Environmental Factors Seasonal Trends – Leaf Nitrogen Concentration



Cultural and Environmental Factors Seasonal Trends – Leaf Potassium and Phosphorus Concentration



How can we quantify environmental and cultural factor effects on plant processes – Photosynthesis?



Temporal trends in crop growth as affected by environment

Quantifying the Effects of Environmental Factors on Photosynthesis

One way to quantify the effects of environmental factors on photosynthesis is to use environmental productivity Index (EPI) concept:

Actual (Photosynthesis) = Potential * Solar Radiation Index * Water Index * Temperature Index * Nutrient Indices (C, N, P, K) * UV-B Index, Salt stress Index, Flooding Index, Ozone Index, etc.,

First, we have to calculate the potential photosynthesis for a given species or cultivar. Potential photosynthesis is defined as the amount of photosynthesis that takes place at a maximum solar radiation under optimum environmental conditions (optimum water, nutrient, zero UV-B, temperature (27 °C) and in an actively growing canopy, no aging effect).

Quantifying the Effects of Environmental Factors on Photosynthesis

Then, we have to account for all the environmental factors that limit to obtain that potential.

Individual environmental factors affect the potential photosynthesis multiplicatively, not additively. For instance, if prolonged drought causes daily stomatal opening to cease, then no photosynthesis will occur, regardless of whether or not light, temperature or other factors are optimal for photosynthesis.

All the indices, ranging from 0 when it is totally limiting photosynthesis to 1 when it does not limit photosynthesis, represent the fractional limitation due to that particular environmental factor. Therefore, photosynthesis decreases as the effect of that particular stress becomes more severe.

Quantifying the Effects of Environmental Factors on Photosynthesis

This way, we could be able to quantify the effect of all environmental factors limiting crop photosynthesis in multi-stress environments or in field conditions.

Quantifying the Effects of Environmental Factors on Photosynthesis

Database and Modeling Methodologies with Cotton as an Example Crop

Crop Responses to Environment - Tools




Soil-Plant-Atmosphere-Research (SPAR) Facility Controlling Environmental Variables

Soil-Plant-Atmosphere-Research (SPAR) Facility

Temperature = 30/22 °C (Average = 27 °C)
and in ambient (360 ppm) CO₂ conditions.

SPAR - Data Acquisition Atmospheric Carbon Dioxide Control



Soil-Plant-Atmosphere-Research (SPAR) Facility

Measuring Gas Exchanges
Carbon [CO₂] Fluxes


Measuring Carbon Fluxes

Carbon Fluxes: Mass balance approach


During sunlit hours, by maintaining steady or constant CO₂ concentration inside the SPAR chamber, we can calculate,

Net photosynthesis = Amount of CO₂ injected – leak rate


Gross Photosynthesis = Net photosynthesis + Respiration




Canopy Photosynthesis Response to Solar Radiation









Estimating Potential Photosynthesis for Cotton as a Function of Solar Radiation



Weather Variables - Mississippi State - 1992 Temporal Trends in Light Interception - 18 May = 0





Materials and Methods


Data collection:
Leaf water potential
(LWP)




Pressure bomb




Canopy Photosynthesis - Environment Response to Water Deficits




Canopy Photosynthesis - Environment Response to Water Deficits




Canopy Photosynthesis - Environment Response to UV-B Radiation






Canopy Photosynthesis - Environment Response to UV-B Radiation



Canopy Photosynthesis - Environment Response to Fertilization - Nitrogen





- ### Photosynthesis and Environment
- Modeling photosynthesis:**
- ✓ Daily values of environmental variables such as temperature and solar radiation (total as well as UV-B) as inputs (Physical inputs).
 - ✓ Daily values of light interception (A separate model for solar radiation interception).
 - ✓ Daily values of leaf nutrient (N,P, K) status (Models for nutrient uptake and leaf nutrient status).
 - ✓ Daily values of leaf water potential as affected by precipitation and irrigation (Model for water uptake and leaf water potential).
 - ✓ Daily values of soil salt concentration (Model for salt concentration).
 - ✓ Soil oxygen concentration (Model for oxygen concentration based on rainfall and irrigation).

Photosynthesis and Respiration and Environment

Actual photosynthesis:

Potential photosynthesis ($159.07 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$)*EPI Indices (solar radiation, Temperature, Water stress, Nutrient stresses, UV-B radiation, salt, and flooding stresses) for various environmental factors.

Therefore, EPI is the way to quantify the effects of environmental factors on photosynthesis and thus productivity of any crop.

- ### Environmental Productivity Index (EPI)
- Same concept can be applied for other crop growth and developmental processes.
 - The EPI concept has universal applicability and NOT location or crop-specific.
 - EPI also allows one to interpret and to understand stresses in the field situations.
 - If we know the factor that is limiting most at any point of time during the growing season, then we can make appropriate management decisions to correct that limitation.

Environmental Productivity Concept Environment - Photosynthesis

Application of Environmental
Productivity Index Concept to the Real-
World Situation


Environmental Factors Impacting Photosynthesis, Productivity and Growth of Crops in a Single Season

Let us examine the environmental variables impacting crop growth and development in a single growing season:

Location: Mississippi State, North Farm
Year : 1992 cotton growing season
Cultivar: DPL 90
Fertilizer Applications: 80 lb N prior to planting
Irrigation/rain-fed: Rain-fed only
Pesticide and weed control: Standard best management practices


Weather Variables - Mississippi State - 1992

Temporal Trends in Temperatures - 18 May = 0




Weather Variables - Mississippi State - 1992

Temporal Trends in Solar Radiation - 18 May = 0




Weather Variables - Mississippi State - 1992

Temporal Trends in Precipitation - 18 May = 0




Weather Variables - Mississippi State - 1992


Temporal Trends in Wind Run - 18 May = 0



Impact of Weather on Plant Growth - Mississippi State - 1992
 Temporal Trends in Mainstem Nodes - Simulated and Observed




Impact of Weather on Plant Growth - Mississippi State - 1992
 Temporal Trends in Plant Height - Simulated and Observed




Quantifying the Effects of Environmental Factors on Photosynthesis

Let us assume the following crop conditions for leaf nitrogen, leaf K, and midday leaf water potential and weather variables such as solar radiation and use percent light interception to calculate an intercepted portion of the incoming solar radiation and temperatures for applying the EPI concept for one cotton growing season - 1992.


Weather Variables - Mississippi State - 1992
 Temporal Trends in Temperatures - 18 May = 0






Weather Variables - Mississippi State - 1992
 Temporal Trends in Solar Radiation - 18 May = 0




Weather Variables - Mississippi State - 1992
 Temporal Trends in Light Interception - 18 May = 0





Applying EPI Concept to Real-world Situation

1. First potential photosynthesis is calculated at optimum temperature, water, and nutrient conditions and 0 UV-B and at maximum solar radiation in an actively growing canopy. **That is equal to 159.07 g CO₂ m² d⁻¹.**
2. Then, using the functional algorithms or equations for Solar radiation, UV-B radiation, temperature, water stress, and nutrient stresses, EPI Indices for the environmental factors are calculated.
3. Finally, actual photosynthesis is estimated = Potential * EPI indices for various environmental factors.



Applying EPI Concept to Real-world Situation

Potential photosynthesis = 159.07 g CO₂ m² d⁻¹ at 25 MJ m² d⁻¹.

Then, actual photosynthesis = potential * EPI-solar radiation * EPI-UV-B radiation * EPI-temp * EPI-CO₂ * EPI-water * EPI-leaf N * EPI-leaf K

Where:

EPI for solar Radiation = 0.0678*intercepted radiation - 0.001111*intercepted radiation²

EPI for Temp = -2.0247 + (0.2141 *Temp) - (0.003779 *Temp²)




EPI for CO₂ = 0.004050 *CO₂ - (0.000004006 *CO₂²) + (0.000000001303*CO₂³)

EPI for Water = 1.3129 + (0.2608 * LWP)

EPI for N = -0.4029 + (0.5954 * Leaf N) - (0.0630 * Leaf N²)

EPI for K = 1.0028 * (1-exp (-1.4577*Leaf K))

EPI for UV-B = 0.9835 - (0.0002563*UV-B) - (0.002163*UV-B²)



Radiation Totals for the 1992 Growing season Mississippi State – North Farm

Variable	Amount, MJ
Total Incoming Radiation	2842
Intercepted Radiation	1551
Percent Intercepted	55

Photosynthesis – EPI Concept Accounting for Individual factors

Variable	Amount, g CO ₂ m ⁻² season ⁻¹
Incoming R	19644
Intercepted R	11441 (100%)
Int. R * UV-B	10448 (9%)
Int. R * T	10139 (11%)
Int. R * W	9783 (14%)
Int. R * N	8986 (21%)
Int. R * K	10841 (5%)

Photosynthesis – EPI Concept Accounting for Multiple Factors

Variable	Amount, g CO ₂ m ⁻² season ⁻¹
Incoming R	19644
Intercepted R	11441 (100%)
Int. R * UV-B	10448 (9%)
Int. R * UV-B * T	9153 (20%)
Int. R * UV-B * T * W	7551 (34%)
Int. R * UV-B * T * W * N	6292 (55%)
Int. R * UV-B * T * W * K	4576 (60%)

Actual amount

Applying EPI Concept to Real-world Situation

1. Here, we have seen the demonstration EPI concept in cotton for the whole growing season to estimate canopy photosynthesis.
2. Potential photosynthesis under optimum conditions; $159.07 \text{ g CO}_2 \text{ m}^{-2} \text{ d}^{-1}$.
3. Then, using the functional algorithms or equations for solar radiation, UV-B radiation, temperature, water stress, and nutrient stresses, and applying EPI indices for various environmental factors to estimate actual photosynthesis.
4. Finally, actual photosynthesis is estimated = Potential *EPI indices for various environmental factors.